

Delayed Against Rapid Sequence Induction in Adults With Medical Diseases Undergoing Emergency Endotracheal Intubation.

NCT06940219

Status	RECRUITING
Phase	Not Applicable
Sponsor	Leipzig University Medical Center
Enrollment	200 participants

Key Eligibility Criteria

Inclusion (2)

- Adult patients 18 years or older in need of emergency endotracheal intubation in the following 60 minutes.
- The planned operator routinely performs endotracheal intubation in critically ill patients.

Exclusion (6)

- Endotracheal Intubation performed during cardiopulmonary resuscitation or during "unstable ROSC", defined as Norepinephrine or Epinephrine dose $> 0,5 \mu\text{g}/\text{kg}/\text{min}$ after return of spontaneous circulation.
- Participation in the study is not warranted because of increased risk for the patient or otherwise not justifiable (e.g. active oral bleeding).
- Known allergy against Ketamine, Esketamine, Midazolam or Rocuronium.
- Known contraindication against Ketamine, Esketamine Midazolam or Rocuronium.
- Known or anticipated difficult airway with indication for awake fiberoptic intubation.

... and 1 more (see full listing online)

Locations (1 total)

Interdisciplinary Medical ICU, Leipzig Medical University Center, Leipzig, Saxony, Germany