

Effects of Cannabidiol on Stress and Nicotine Withdrawal

NCT07001930

Status	RECRUITING
Phase	Phase 1
Sponsor	Johns Hopkins University
Enrollment	90 participants

Key Eligibility Criteria

Inclusion (6)

- Healthy non-treatment seeking adults aged 21-70 (inclusive).
 - Use of ≥ 5 cigarettes per day for ≥ 2 years. (Use of other tobacco/nicotine products in addition to cigarettes will be acceptable).
 - Biological confirmation of cigarette use: have an expired carbon monoxide (CO) level ≥ 8 ppm.
 - Alcohol Use: < 2 drinks/day on average (< 14 drinks/week).
 - Sleep schedule: No graveyard or third shift work (i.e., participants must report a normal nighttime sleep schedule).
- ... and 1 more (see full listing online)

Exclusion (11)

- Report current intention to reduce or quit cannabis or tobacco use within the next 30 days
 - Meet Diagnostic and Statistical Manual (DSM-5) criteria for a substance use disorder other than alcohol, cannabis, or nicotine
 - Test positive for illicit drugs other than cannabis
 - Positive breath alcohol test ($\geq .01$ g/210L) at study admission
 - Have a current physical or mental illness or takes a medication judged by the study team to negatively impact participant safety or scientific integrity
- ... and 6 more (see full listing online)

Locations (1 total)

Behavioral Pharmacology Research Unit, Baltimore, Maryland, United States