

Non-inferiority Trial Comparing Vaginal Ultrasound-guided Ethanol Sclerotherapy and Laparoscopic Cystectomy for the Management of Ovarian Endometriomas (E-STEP)

NCT07004959

Status	RECRUITING
Phase	Not Applicable
Sponsor	Assistance Publique Hopitaux De Marseille
Enrollment	64 participants

Key Eligibility Criteria

Inclusion (3)

- Patients aged between 18 and 45
- Endometrioma diagnosed by endovaginal ultrasound or MRI (examinations carried out by a radiologist or gynaecologist) with a typical appearance and whose largest diameter is between 3 and 10 centimetres.
- Pain (dysmenorrhoea or chronic pelvic pain) with a VAS greater than 30 millimetres, assessed by the question "Indicate the subjective level of your endometriosis-related pain over the last 4 weeks".

Exclusion (10)

- Inability to speak French
 - Refusal to take part in the study
 - Risk factors for impaired ovarian reserve:
 - Ovarian reserve already low (AMH $\leq 1,5$ ng/ml or Antral follicle count ≤ 7 on the contralateral ovary)
 - Bilateral or multiple endometriomas (two or more endometriomas greater than 3 centimetres in diameter).
- ... and 5 more (see full listing online)

Locations (1 total)

gynecology department APHM, Marseille, France, France