

# Strategies for Antithrombotic Treatment Following Transcatheter Edge-to-Edge Repair in Patients Without an Indication for Oral Anticoagulant

NCT07007143

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 4
<b>Sponsor</b>	Chinese Academy of Medical Sciences, Fuwai Hospital
<b>Enrollment</b>	1,032 participants

## Key Eligibility Criteria

---

### Inclusion (5)

- Successful TEER procedure, defined as technical success per MVARC criteria.
- Ability and willingness to comply with the trial protocol.
- Provision of written informed consent.
- Women of childbearing potential must use effective contraception from the time of consent until the final dose of antithrombotic therapy.
- Antithrombotic strategy approved by the investigator.

### Exclusion (13)

- Severe renal impairment (creatinine clearance  $< 15$  mL/min or requiring dialysis).
- Ongoing postoperative bleeding (defined as overt bleeding with a  $\geq 3.0$  g/dL drop in hemoglobin or requiring  $\geq 3$  units of blood transfusion), or vascular complications following the index TEER procedure.
- Platelet count  $< 50 \times 10^9$  /L.
- Need for reoperation due to complications of the index TEER procedure.
- Recent ( $< 12$  month) intracranial or intracerebral hemorrhage.

... and 8 more (see full listing online)

## Locations (1 total)

---

Chinese Academy of Medical Sciences, Fuwai Hospital, Beijing, Beijing Municipality, China