

ICF-Based Effects of PNF and Kinesio Taping on Hand Function in Hemiplegic Stroke

NCT07014696

Status	RECRUITING
Phase	Not Applicable
Sponsor	Istinye University
Enrollment	54 participants

Key Eligibility Criteria

Inclusion (9)

- Stroke diagnosed by MRI or CT, regardless of ischemic or hemorrhagic origin,
- Hemiplegia caused by stroke alone,
- At least 6 and at most 24 months have passed since the stroke was diagnosed
- Being over 18 years of age,
- Volunteering to participate in the study

... and 4 more (see full listing online)

Exclusion (9)

- Individuals with bilateral hemiplegia
- Individuals with unstable vital signs
- Individuals with any open wound in the upper extremity
- Pathological conditions affecting upper extremity sensation (undergoing surgery after fracture, etc.)
- Individuals with a BMI value greater than 29.9 kg/m²

... and 4 more (see full listing online)

Locations (2 total)

Ostinye University, Bahçe_ehir Liv Hospital, Istanbul, Bahçe_ir, Turkey (Türkiye)

Ondokuz Mayıs Üniversitesi, SUVAM Havza Fizik Tedavi ve Rehabilitasyon Merkezin, Samsun, Havza, Turkey (Türkiye)