

Randomized Study on Transoral Outlet Reduction and Lifestyle Changes for Weight Regain After Gastric Bypass

NCT07015424

Status	RECRUITING
Phase	Not Applicable
Sponsor	Boston Scientific Corporation
Enrollment	108 participants

Key Eligibility Criteria

Inclusion (12)

- years or older.
- Experiencing weight regain (e 20% weight from nadir) after achieving clinically successful weight loss.
- Willing to have a TORe procedure for the treatment of weight regain.
- BMI between 30 and 50 kg/m2, inclusive.
- GJA diameter e 20 mm, confirmed either during a

... and 7 more (see full listing online)

Exclusion (6)

- Current use or use within the three months prior to the baseline visit of over the counter or prescribed weight loss supplements / medications (including incretins).
- Current or planned management of Type 2 Diabetes with incretin medications.
- Inability or unwillingness to comply with assigned treatment or protocol requirements related to visits.
- Investigator's medical judgement that the subject is not a candidate for the TORe procedure.
- Current pregnancy confirmed by site standard of care or planned pregnancy or breastfeeding in the next two years.

... and 1 more (see full listing online)

Locations (9 total)

Hoag Memorial Hospital, Irvine, California, United States
Stanford University Medical Center, Stanford, California, United States
Mayo Clinic, Jacksonville, Florida, United States
... and 6 more locations