

Comparison of the Effectiveness of Paracetamol With Ibuprofen or Paracetamol With Metamizole in Treating Pain in Acute Pancreatitis in Children

NCT07024199

Status	RECRUITING
Phase	Phase 4
Sponsor	Medical University of Warsaw
Enrollment	78 participants

Key Eligibility Criteria

Inclusion (5)

- diagnosis of AP according to the INSPPIRE mentioned above criteria,
- age from 3 to 18 years of age,
- abdominal pain on admission assessed on the Numerical Rating Scale (NRS) or FLACC ≥ 4 points,
- no analgesic treatment before enrolment in the study OR the last dose of analgesic drug (paracetamol, ibuprofen, metamizole) taken ≥ 6 hours before enrolment for examination,
- consent of legal guardians and the child (in the case of patients ≥ 16 years of age) to participate in the study.

Exclusion (12)

- who took the last dose of painkiller (paracetamol, ibuprofen, metamizole) < 6 hours before entering the study,
- allergic to acetylsalicylic acid, other NSAIDs, paracetamol, metamizole,
- with inflammatory bowel disease,
- with gastrointestinal bleeding and other active bleeding,
- with gastric and/or duodenal ulcer disease,
- ... and 7 more (see full listing online)

Locations (1 total)

Department of Paediatric Gastroenterology and Nutrition, Warsaw, Poland