

# Effectiveness of Speech Therapy for Lip-reading in Patients With Hearing Aids: SCED Study

NCT07028658

---

**Status** RECRUITING  
**Sponsor** Institut Pasteur  
**Enrollment** 8 participants

## Key Eligibility Criteria

---

### Inclusion (8)

- Patient aged 18 or older
- Affiliated with a social security system
- Moderate to severe bilateral post-lingual hearing loss, fitted with hearing aids for more than 6 months
- Wearing hearing aids for a minimum of 8 hours/day on average over the past month
- Intelligibility in quiet  $\geq 80\%$  correct responses at 60 dB (Lafon cochlear list)
- ... and 3 more (see full listing online)

### Exclusion (5)

- Patient who has already undergone speech therapy rehabilitation as part of their hearing loss treatment
- Diagnosis of neurodegenerative disease or history of brain injury (traumatic brain injury, stroke, tumor, meningitis, etc.)
- Uncorrected vision or visual impairment affecting lip reading (e.g. AMD, glaucoma, unoperated cataract, etc.)
- Presence of severe tinnitus (score  $\geq 56$ ) on the Tinnitus Handicap Inventory (THI) questionnaire
- Significant cognitive impairment (CODEX C or D) or Montreal Cognitive Assessment (MOCA) score  $\leq 23$

## Locations (1 total)

---

Cabinet d'orthophonie, Paris, France