

The Effect of Systemic Nonsteroidal Anti-inflammatory Drugs (NSAIDs) Vs Intraoperative Infiltration of Steroids in Tonsillar Bed Following Tonsillectomy on Post Tonsillectomy Pain.

NCT07035314

Status	RECRUITING
Phase	Not Applicable
Sponsor	Ain Shams University
Enrollment	75 participants

Key Eligibility Criteria

Inclusion (6)

- Age: Between 4-10 years.
- Indications of tonsillectomy such as:
- Recurrent acute attacks at least 7 episodes in the past year or at least 5 episodes per year for 2 years or at least 3 episodes per year for 3 years.
- Chronic tonsillitis and hypertrophy of tonsils causing sleep apnea, difficulty in deglutition, interference in speech.
- Tonsillectomy by cold dissection only.
- ... and 1 more (see full listing online)

Exclusion (7)

- Patients with age less than 4 and more than 10 years.
- Indications of tonsillectomy other than chronic tonsillitis such as lymphoma.
- Patients on chronic steroid therapy
- Hemoglobin level less than 10 gm/dL
- Presence of acute infection in the upper respiratory tract, acute tonsillitis
- ... and 2 more (see full listing online)

Locations (1 total)

Ainshams university hospitals, Cairo, Egypt

<https://clinicaltrials.gov/study/NCT07035314>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).