

# Comparison of Distalization of Maxillary First and Second Molars Between a Miniscrew Implant Supported Device and Clear Aligners.

NCT07040176

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Aristotle University Of Thessaloniki
Enrollment	40 participants

## Key Eligibility Criteria

---

### Inclusion (3)

- Patients with at least half cusp bilateral Class II malocclusion between upper and lower first molars due to maxillary excess or maxillary protrusion or forward movement of the maxillary posterior teeth, where maxillary molar distalization is indicated.
- Patients in the permanent dentition period of less than 30 years of age with second molars erupted at least one half.
- Patients with good general health and healthy periodontium.

### Exclusion (7)

- Patients with Class II malocclusion between upper and lower first molars due to mandibular deficiency or mandibular retrusion.
- Patients with less than half cusp bilateral Class II malocclusion between upper and lower first molars.
- Patients with unilateral Class II malocclusion between upper and lower first molars.
- Patients with cleft lip and palate or with other craniofacial syndromes.
- Patients with missing, or congenital missing or supernumerary teeth.

... and 2 more (see full listing online)

## Locations (1 total)

---

Department of Orthodontics, Faculty of Dentistry, School of Medical Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece