

Safety and Efficacy Evaluation of LX111 Gene Therapy in nAMD Patients

NCT07053358

Status	RECRUITING
Phase	Early Phase 1
Sponsor	Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine
Enrollment	10 participants

Key Eligibility Criteria

Inclusion (6)

- Willing to sign the informed consent, and willing to attend follow-up visits.
- Age \geq 50
- Diagnosis of active CNV secondary to neovascular AMD
- BCVA ETDRS letters between 5 and 63
- Subjects must have received a minimum of 2 injections within 6 months prior to screening and demonstrated a meaningful response to anti-VEGF therapy
- ... and 1 more (see full listing online)

Exclusion (5)

- CNV or macular edema in the study eye secondary to diseases other than nAMD
- Retinal detachment, uveitis, uncontrolled glaucoma in the study eye, or any condition preventing visual acuity improvement
- Acute coronary syndrome, myocardial infarction or coronary artery revascularization, CVA, TIA in the last 6 months
- Uncontrolled hypertension defined as average SBP \geq 160 mmHg or an average DBP \geq 100 mmHg
- Uncontrolled diabetes defined as HbA1c \geq 8.0% within 28 days prior to screening

Locations (1 total)

Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China