

# Food Effect on PK of DW-1021 (Pelubiprofen 45 mg / Tramadol 45.9 mg) in Healthy Adults

NCT07060209

---

Status	RECRUITING
Phase	Phase 1
Sponsor	Haiphong University of Medicine and Pharmacy
Enrollment	14 participants

## Key Eligibility Criteria

---

### Inclusion (9)

- Healthy male subjects aged 20 to 40 years at screening visit
- Body Mass Index (BMI) between 18.5 and 24.9 kg/m<sup>2</sup>
- Body weight greater than 50 kg
- Systolic blood pressure between 100 mmHg and 129 mmHg; diastolic blood pressure less than 84 mmHg
- Regular heart rate ranging from 60 to 90 beats per minute

... and 4 more (see full listing online)

### Exclusion (8)

- Use of drugs that induce or inhibit drug-metabolizing enzymes (e.g., barbiturates) within 30 days prior to administration, or use of any medication that might affect the study within 10 days prior to administration
- Participation in any other clinical trial within 3 months prior to screening
- Blood donation within 8 weeks prior to drug administration
- History of gastrointestinal surgery that may affect drug absorption
- History of drug abuse, or use of alcohol, drugs, or tobacco products within 1 year before participation

... and 3 more (see full listing online)

## Locations (2 total)

---

Clinical Trial and Bioequivalence Center, Haiphong, Hai Phong, Vietnam  
Clinical Trial and Bioequivalence Center, Haiphong, Hai Phong, Vietnam