

Effects of Frontopolar TMS in Alcohol Craving

NCT07064590

Status	RECRUITING
Phase	Not Applicable
Sponsor	Goethe University
Enrollment	34 participants

Key Eligibility Criteria

Inclusion (4)

- adults aged 18-65
- ICD-10 diagnosis of alcohol dependence
- Ability to give consent
- Sinus rhythm in ECG

Exclusion (4)

- Current psychotic symptoms in patients with psychotic disorders (F20, F23, F10.5)
- Contraindication against TMS or MRI
- Severe neurological disorders such as epilepsy, stroke, neuroinflammatory disorders (e.g., multiple sclerosis). A history of seizures only in the context of alcohol withdrawal does not represent a contraindication, unless the seizure happened in the last 3 weeks before study inclusion.
- Acute withdrawal symptoms (CIWA-Ar \gt 5)

Locations (2 total)

Klinik und Poliklinik für Psychiatrie, Psychosomatik und Psychotherapie Universitätsklinikum Frankfurt, Goethe-Universität, Frankfurt am Main, Hesse, Germany
Klinik und Poliklinik für Psychiatrie, Psychosomatik und Psychotherapie Universitätsklinikum Frankfurt, Goethe-Universität, Frankfurt am Main, Hesse, Germany