

Effect of Lymphovenous Anastomosis on Lower Limb Lymphedema: Pragmatic Randomized Controlled Trial

NCT07073586

Status	RECRUITING
Phase	Not Applicable
Sponsor	Herlev and Gentofte Hospital
Enrollment	66 participants

Key Eligibility Criteria

Inclusion (6)

- LLL International Society of Lymphology (ISL) stage 1-2a (both primary and secondary lymphedema and both unilateral and bilateral)
 - Cancer-free with no recurrence for at least one year
 - Age 18 or above
 - Circumference of the affected leg is at least 1 cm larger than the non-lymphedema leg at the most affected site of lymphedema.
 - Proficiency in Danish language, and ability to provide informed consent.
- ... and 1 more (see full listing online)

Exclusion (14)

- LLL International Society of Lymphology (ISL) stage 2b-3
 - Medical conditions contraindicating surgical intervention or anesthesia, such as severe heart or lung disease
 - Allergy to ICG
 - Ongoing infections or skin diseases in the affected limb
 - Previous surgery or other treatment modalities that could interfere with the study results (previous LVA, liposuction or similar).
- ... and 9 more (see full listing online)

Locations (2 total)

Department of Plastic Surgery, Odense University Hospital, Odense, Fyn, Denmark
Department of Plastic Surgery, Herlev and Gentofte Hospital, Herlev, Region Sjælland, Denmark