

# An Open-label, Multicenter Study to Assess the Pharmacokinetics (PK), Safety, and Tolerability of Subcutaneous IgPro20 in Immunoglobulin (IG) Treatment-naïve Participants With Primary Immunodeficiency (PID)

NCT07076446

---

Status	RECRUITING
Phase	Phase 4
Sponsor	CSL Behring
Enrollment	8 participants

## Key Eligibility Criteria

---

### Inclusion (2)

- Participants must be aged  $\geq 18$  years.
- Participants must have a confirmed and documented diagnosis of PID, must be IG treatment-naïve and have an IgG level less than or equal to ( $\leq$ ) 400 milligrams per deciliter (mg/dL) at Screening.

### Exclusion (13)

- Participants with hyperprolinemia.
- Participants who are receiving the following medications:
  - Systemic corticosteroids (prednisone or equivalent; average daily dose of greater than  $> 15$  mg) from 4 weeks before Screening.
  - Any dose of systemic immunosuppressants within 9 months or 5 times the half-life ( $t_{1/2}$ ) plus 6 months before Screening, whichever is longer.
  - Any dose of biologic therapies with influence on the immune system (eg, tumor necrosis factor inhibitors, interleukin inhibitors, B-cell inhibitors), including investigational agents, within 12 months or 5 times the  $t_{1/2}$  plus 6 months before Screening, whichever is longer.
- ... and 8 more (see full listing online)

## Locations (8 total)

---

Research Solutions of AZ, Litchfield Park, Arizona, United States  
Medical Research of Arizona, Scottsdale, Arizona, United States  
Immuno Health Centers, Centennial, Colorado, United States  
... and 5 more locations