

Safety and Performance, and Dose, of OsteoAdapt DE in Localized Alveolar Ridge Augmentations

NCT07078331

Status	RECRUITING
Phase	Phase 1, Phase 2
Sponsor	Theradaptive, Inc
Enrollment	45 participants

Key Eligibility Criteria

Inclusion (8)

- Age 22 or older at the time of signing informed consent
 - Subject requires extraction of a single tooth due to extensive carious lesions, prosthetic or endodontic failure, root fracture, or other reasons with dental implant treatment planned at the site
 - Tooth root in position allowing a bone core sample to be harvested within the former socket
 - mm of alveolar bone height without impinging on the maxillary sinus or inferior alveolar canal
 - Dehiscence of d 2 mm on the buccal bony wall at the time of extraction
- ... and 3 more (see full listing online)

Exclusion (16)

- Active localized or systemic infection
 - Untreated periodontal disease
 - Presence of a fenestration larger than 3 mm in diameter in the buccal wall at the time of extraction
 - Inadequate bone dimensions or restorative space to place a dental implant
 - The subject uses, or has used within 30 45 days of surgery, tobacco or nicotine or is prescribed steroids such as cortisone
- ... and 11 more (see full listing online)

Locations (6 total)

Armenia Medical Center, Yerevan, Armenia
Medline Clinic, Yerevan, Armenia
Hadassah Medical Center, Jerusalem, Israel
... and 3 more locations