

Short-term Bactericidal Effect of Contezolid in MAC-PD

NCT07084402

Status	RECRUITING
Phase	Phase 4
Sponsor	Bin Cao
Enrollment	188 participants

Key Eligibility Criteria

Inclusion (12)

- Patients voluntarily participate in this study and sign the Informed Consent Form.
 - Age e 18 years and d75 years; gender unrestricted.
 - Confirmed diagnosis of MAC pulmonary disease per ATS/IDSA 2020 guidelines or Chinese Guidelines for Diagnosis and Treatment of Nontuberculous Mycobacterial Diseases (2020 edition), with imaging features consistent with nodular bronchiectatic type.
 - No prior anti-MAC treatment within the 3 months preceding screening.
 - For premenopausal women of childbearing potential who are not surgically sterile:
- ... and 7 more (see full listing online)

Exclusion (13)

- History of allergy to any study drug in the treatment regimen.
 - Mixed infections with multiple mycobacteria, bacteria, fungi, or viruses (e.g., HIV coinfection).
 - Presence of congenital/acquired immunodeficiency disorders, active pulmonary malignancy (primary or metastatic), or other malignancies requiring chemotherapy/radiation therapy during the screening or study period.
 - History of solid organ transplantation.
 - Currently undergoing dialysis.
- ... and 8 more (see full listing online)

Locations (3 total)

Beijing Chest Hospital, Capital Medical University, Beijing, Beijing Municipality, China
China-Japan Friendship Hospital, Beijing, Beijing Municipality, China
Jiangxi Provincial Chest Hospital, Nanchang, Jiangxi, China