

# Clinical Effects of Keratinized Mucosa Augmentation Around Dental Implant With Free Gingival Graft in Different Surgical Timing at Mandibular Posterior Sites

NCT07088172

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	The Dental Hospital of Zhejiang University School of Medicine
<b>Enrollment</b>	24 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Age e 18 years, gender unrestricted;
- partially edentulous in the mandible posterior area, planned for implant-supported fixed restoration;
- Buccal keratinized mucosa width at the implant site is less than 2mm, with the transplantation area not exceeding 2 teeth;
- No soft tissue augmentation performed at the site prior to surgery;
- Overall health is good, with no systemic diseases that may affect bone metabolism or wound healing;
- ... and 2 more (see full listing online)

### Exclusion (5)

- Uncontrolled metabolic diseases;
- Severe periodontal disease, with unstable periodontal condition;
- Smoking \> 10 cigarettes per day;
- Unwilling to participate in the follow-up of this study;
- Other conditions that do not meet the requirements for implant treatment.

## Locations (1 total)

---

The Stomatology Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China