

# Lacunar Stroke hyperAcute Clinical Utilization of Novel Approach Regimens: Rt-PA vs. DAPT Randomised Clinical Trial

NCT07111559

---

Status	RECRUITING
Phase	Phase 4
Sponsor	Nippon Medical School
Enrollment	500 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Age e 18 years.
- Acute ischemic stroke within 4.5 hours from onset. If onset time is unknown because of impaired consciousness or aphasia, use the "last known well" time.
- A single perforating-artery infarct on brain MRI:
  - located in the corona radiata, putamen, internal capsule, thalamus, or pons; solitary, mainly round or oval, with a maximum diameter d 20 mm; lesions only in the centrum semiovale are not allowed, but extension from the above sites into the centrum semiovale is allowed.
- No disability in daily life before the stroke (modified Rankin Scale d 1).

... and 2 more (see full listing online)

### Exclusion (7)

- Antithrombotic therapy considered inappropriate because of active bleeding, low platelet count, or similar conditions.
- Any contraindication to intravenous rt-PA, without blood pressures.
- e 50 %stenosis or occlusion of the artery responsible for the stroke \\* (see note below).
- Diseases that require anticoagulation (e.g., atrial fibrillation, deep-vein thrombosis) \\*
- Inability to take medicine orally.

... and 2 more (see full listing online)

## Locations (27 total)

---

University of Yamanashi Hospital, ChkM, Japan  
Fukuoka Red Cross Hospital, Fukuoka, Japan  
Ota Memorial Hospital, Fukuyama, Japan

... and 24 more locations