

Effect of Collagen Membranes on Distal Periodontal Healing of Second Molars After Impacted Mandibular Third Molar Surgery.

NCT07114523

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Medicine and Pharmacy at Ho Chi Minh City
Enrollment	20 participants

Key Eligibility Criteria

Inclusion (5)

- Patients aged 25 years or older.
- Patients with mesioangular or horizontally impacted mandibular third molars. The highest point of mandibular third molars are lower than the occlusal surface of adjacent second molars (classified B, C according to Pell - Gregory's category).
- Patients with symmetrical lower third molars on both sides with same classification according to Pell - Gregory's category. The difference of third molars inclination on both side is not more than 15°.
- Patients with lower second molars on both sides have no cavities or fillings that affect the determination of the cemento-enamel junction.
- Patients are explained, counseled clearly and sign their consent to participate in the study.

Exclusion (8)

- Patients with coagulation or immune disorders.
- Patients allergic to local anesthetics (Lidocaine 2% with Adrenaline 1:100,000) or any drugs used in the study.
- Patients allergic to products or food derived from pigs.
- Pregnant or breastfeeding patients.
- Patients who smoke.

... and 3 more (see full listing online)

Locations (1 total)

University of Medicine and Pharmacy at Ho Chi Minh city, Ho Chi Minh City, Vietnam