

Clinical Evaluation of With or Without Internal Limiting Membrane Peeling for Idiopathic Epiretinal Membrane Surgery: A Randomized Controlled Trial

NCT07142785

Status	RECRUITING
Phase	Not Applicable
Sponsor	Second Affiliated Hospital, School of Medicine, Zhejiang University
Enrollment	96 participants

Key Eligibility Criteria

Inclusion (3)

- Patients aged ≥ 18 years presenting to our hospital within the study period.
- Patients diagnosed with idiopathic epiretinal membrane (ERM) based on examination.
- Patients with best-corrected visual acuity (BCVA) ≤ 0.6 or the presence of metamorphopsia.

Exclusion (6)

- Patients with secondary ERM identified during preoperative evaluation (e.g., secondary to retinal vein occlusion).
- Patients with concomitant ocular diseases affecting the macula, such as macular degeneration, macular schisis, or macular atrophy.
- Patients with a history of ocular trauma or previous ocular surgery (except for cataract surgery).
- Patients with severe systemic diseases contraindicated for surgery.
- Patients with poor compliance or concomitant psychiatric disorders.

... and 1 more (see full listing online)

Locations (1 total)

Eye Center of the Second Affiliated Hospital of Zhejiang University, Hangzhou, Zhejiang, China