

The Efficacy and Safety of Pregabalin and Mirogabalin in Patients With Fibromyalgia

NCT07157852

Status	RECRUITING
Phase	Not Applicable
Sponsor	Beijing Tiantan Hospital
Enrollment	674 participants

Key Eligibility Criteria

Inclusion (7)

- Newly diagnosed with FM according to the 2016 Revisions to the 2010/2011 FM diagnostic criteria;\[1\]
- Aged over 18 years old;
- Suffering from moderate to severe FM, refractory to non-pharmacological interventions and without prior exposure to recommend pharmacological treatments for FM;
- Baseline numeric rating scale (NRS) score of 4 or higher;
- Aspartate aminotransferase and alanine aminotransferase levels below twice the upper limit of normal range;
- ... and 2 more (see full listing online)

Exclusion (5)

- Previous allergic reactions to pregabalin, mirogabalin, or any of their excipients;
- Prior diagnosis of epilepsy or depression requiring antidepressant therapy;
- Women who are pregnant or breastfeeding;
- Has severe systemic illnesses, such as poorly controlled hypertension, poorly controlled diabetes mellitus, or significant cardiac impairment;
- Suffering from acute or chronic pain disorders other than FM.

Locations (1 total)

Beijing Tiantan Hospital, Beijing, Beijing 100070, Beijing, China