

Gelaspan vs Crystalloid Therapy in Sepsis

NCT07172451

Status	RECRUITING
Phase	Phase 4
Sponsor	Shanghai Zhongshan Hospital
Enrollment	318 participants

Key Eligibility Criteria

Inclusion (4)

- Age \geq 18 years.
- Diagnosis of sepsis according to the Sepsis-3 definition, caused by intra-abdominal infection, and requiring emergency abdominal surgery for source control.
- Sequential Organ Failure Assessment (SOFA-2) score \geq 2 (determined by the most recent available clinical data) and blood lactate $>$ 2 mmol/L (measured within 6 hours prior to enrollment).
- Subject or legal representative can understand the study purpose and provide written informed consent.

Exclusion (13)

- Received any dose of artificial colloids within 24 hours prior to randomization.
- Predicted mortality within 48 hours (ASA physical status class \geq V).
- Atrial fibrillation or congestive heart failure.
- Severe Acute Respiratory Distress Syndrome (ARDS).
- Preoperative coagulation dysfunction or receiving anticoagulant therapy.

... and 8 more (see full listing online)

Locations (9 total)

The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China

Zhongda Hospital, Southeast University, Nanjing, Jiangsu, China

The First Affiliated Hospital of Soochow University, Suzhou, Jiangsu, China

... and 6 more locations