

# Holy Basil in The Treatment of Dyspepsia

NCT07175272

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 2, Phase 3
<b>Sponsor</b>	Mahidol University
<b>Enrollment</b>	27 participants

## Key Eligibility Criteria

---

### Inclusion (2)

- Participants aged 18 years and older.
- Presence of dyspeptic symptoms, assessed by the Leeds Dyspepsia Questionnaire with a score of at least 5 or higher.

### Exclusion (8)

- Presence of Helicobacter pylori infection.
- Diagnosis of peptic ulcer or erosive esophagitis Los Angeles Classification grade B or greater.
- Use of acid-suppressant therapy including proton pump inhibitors (PPIs) or histamine-2 receptor antagonists (H2RAs) that cannot be discontinued at least 2 weeks prior enrolment and during the study period.
- Use of mucosal protective agents, such as rebamipide, sucralfate, or irsogladine that cannot be discontinued at least 4 weeks prior enrolment and during the study period.
- History of gastric cancer or duodenal cancer.

... and 3 more (see full listing online)

## Locations (1 total)

---

Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok Noi, Bangkok, Thailand