

# Comparing Efficacy of Bromfenac 0.09%, Nepafenac 0.3% and Diclofenac 0.1% in Patients After Cataract Surgery

NCT07178639

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Nemocnice Kolín
Enrollment	150 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- Are male or female at least 18 years of age who require cataract surgery and no other surgical procedures during the cataract surgery.
- Agree not to have any other ocular surgical procedures in the study or fellow (non study) eye within 15 days prior to the initiation of dosing with the test article or throughout the duration of the study.
- Have a Best Corrected Visual Acuity of 20/200 or better in either eye.
- Are able to self administer test article (or have a caregiver available to instill all doses of test article).

### Exclusion (15)

- Have known hypersensitivity to bromfenac, nepafenac, diclofenac or any component of the test article (including "procedural" medications such as anesthetic and/or fluorescein drops, dilating drops, etc.).
- Have intraocular inflammation (i.e., cells or flare in the anterior chamber as measured on slit lamp examination) in study eye at screening visit.
- Have a known blood dyscrasia or bone marrow suppression, a diagnosis of uncontrolled/unstable peptic ulcer disease, inflammatory bowel disease, or ulcerative colitis, or any uncontrolled/unstable pulmonary, cardiac, vascular, autoimmune, hepatic, renal, or central nervous system disease.
- Have used ocular, topical, or systemic NSAIDs or ocular, topical, or systemic gentamicin, or cyclosporine ophthalmic emulsion within 7 days prior to initiation of dosing with the test article or throughout the duration of study, with exception of allowing patients on a stable dose of aspirin 81 mg daily or less.
- Have diabetes of any type.

... and 10 more (see full listing online)

## Locations (1 total)

---

Oblastní nemocnice Kolín a.s., Kolín, Czechia