

Time to Ambulation After ProGlide Closure

NCT07179536

Status	RECRUITING
Phase	Not Applicable
Sponsor	Chinese Academy of Medical Sciences, Fuwai Hospital
Enrollment	300 participants

Key Eligibility Criteria

Inclusion (6)

- Scheduled for peripheral angiography or endovascular intervention via transfemoral access.
- Femoral artery puncture site located at the common femoral artery.
- Use of 6-8F vascular sheath.
- Hemostasis achieved with 6F ProGlide closure device (defined as no active bleeding, no hematoma formation, and no ischemia of the punctured limb after closure).
- Preoperative ankle-brachial index (ABI) > 0.9 on both sides.

... and 1 more (see full listing online)

Exclusion (6)

- Undergoing carotid artery intervention.
- Femoral artery diameter < 5 mm, or effective lumen < 5 mm due to plaque burden.
- History of vascular complications at the puncture site.
- Abnormal cardiopulmonary function.
- Intraoperative platelet count $< 80 \times 10^9/L$, or use of thrombolytic agents.

... and 1 more (see full listing online)

Locations (1 total)

Fuwai Hospital, National Center for Cardiovascular Disease, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, Beijing Municipality, China