

The Efficacy and Safety of Pregabalin and Crisugabalin in Patients With Fibromyalgia

NCT07196657

| | |
|------------|--------------------------|
| Status | RECRUITING |
| Phase | Not Applicable |
| Sponsor | Beijing Tiantan Hospital |
| Enrollment | 1,116 participants |

Key Eligibility Criteria

Inclusion (7)

- Diagnosed with FM according to the 2016 Revisions to the 2010/2011 FM diagnostic criteria;
- Aged over 18 years old;
- Suffering from moderate to severe FM, refractory to non-pharmacological interventions and without prior exposure to recommend pharmacological treatments for FM;
- Baseline numeric rating scale (NRS) score of 4 or higher;
- Aspartate aminotransferase and alanine aminotransferase levels below twice the upper limit of normal range;
- ... and 2 more (see full listing online)

Exclusion (5)

- Previous allergic reactions to pregabalin, crisugabalin, or any of their excipients;
- Prior diagnosis of epilepsy or depression requiring antidepressant therapy;
- Women who are pregnant or breastfeeding;
- Has severe systemic illnesses, such as poorly controlled hypertension, poorly controlled diabetes mellitus, or significant cardiac impairment;
- Suffering from acute or chronic pain disorders other than FM.

Locations (1 total)

Beijing Tiantan Hospital, Beijing, Beijing 100070, Beijing, China