

Effects of Early Physiotherapy on Motor Optimality Score in At-Risk of Infants

NCT07201805

Status	RECRUITING
Phase	Not Applicable
Sponsor	Kahramanmaras Sutcu Imam University
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (3)

- Infants diagnosed with periventricular hemorrhage (PVH), intracranial hemorrhage (ICH), cystic PVL, HIE, kernicterus, perinatal asphyxia, neonatal sepsis, necrotizing enterocolitis (NEC), retinopathy of prematurity (ROP), RDS, or BPD, and those receiving oxygen or mechanical ventilation (MV) support
- Infants with a 5-minute Apgar score ≤ 3 , ≤ 37 weeks' gestation, ≤ 1500 grams of preterm birth, or prematurity due to multiple births.
- Infants with the corrected age of 2 to 4 months

Exclusion (1)

- Infants with congenital malformations (spina bifida, congenital muscular torticollis, arthrogryposis multiplex congenita, etc.), babies diagnosed with metabolic and genetic diseases (down syndrome, spinal muscular atrophy, duchenne muscular dystrophy, etc.)

Locations (1 total)

Kahramanmara_ Sütçü imam University, Kahramanmara_, Oniki_übet, Turkey (Türkiye)