

The Efficacy and Safety of Pregabalin Combined With Toludesvenlafaxine in Patients With Fibromyalgia

NCT07208357

Status	RECRUITING
Phase	Not Applicable
Sponsor	Beijing Tiantan Hospital
Enrollment	384 participants

Key Eligibility Criteria

Inclusion (7)

- Diagnosed with FM according to the 2016 Revisions to the 2010/2011 FM diagnostic criteria;
- Aged 18 years or older;
- Experiencing moderate to severe FM that have not been effectively alleviated by non-pharmacological treatments and has not received currently recommended pharmacological treatment for FM;
- Numeric rating scale (NRS) score ≥ 4 at baseline;
- Aspartate aminotransferase and alanine aminotransferase levels less than twice the upper limit of normal;
- ... and 2 more (see full listing online)

Exclusion (5)

- History of hypersensitivity to pregabalin, toludesvenlafaxine or any of its excipients;
- History of epilepsy, or depression requiring antidepressant medications;
- Pregnancy or breastfeeding;
- Presence of serious systemic diseases, including uncontrolled hypertension, uncontrolled diabetes, or significant cardiac dysfunction;
- With acute or chronic pain conditions other than FM.

Locations (1 total)

Beijing Tiantan Hospital, Beijing, Beijing 100070, Beijing, China