

Detrusor Nerve Radiofrequency Ablation for Overactive Bladder in Women

NCT07209397

Status	RECRUITING
Phase	Not Applicable
Sponsor	InMode MD Ltd.
Enrollment	202 participants

Key Eligibility Criteria

Inclusion (9)

- Female, aged 22 to 80 years inclusive at the time of consent.
- History of idiopathic overactive bladder (OAB) symptoms for e6 months.
- At least 7 episodes of urge urinary incontinence (UUI) over 3 days, as recorded in a 3-day bladder diary during screening.
- Urge-predominant incontinence, defined as greater number of UUI episodes than stress urinary incontinence (SUI) episodes during screening.
- Willing and able to complete the 3-day bladder diary.

... and 4 more (see full listing online)

Exclusion (15)

- Predominant stress urinary incontinence (SUI) based on bladder diary
- Currently pregnant, breastfeeding, or planning pregnancy during the study
- History of neurologic conditions affecting bladder function (e.g., multiple sclerosis, spinal cord injury, Parkinson's disease)
- Active urinary tract infection (UTI) at screening
- History of interstitial cystitis, bladder pain syndrome, or chronic pelvic pain

... and 10 more (see full listing online)

Locations (9 total)

Stanford Medicine Department of Obstetrics & Gynecology, Palo Alto, California, United States

MedStar Health Research Institute, Washington D.C., District of Columbia, United States

Advanced Specialty Research, Boise, Idaho, United States

... and 6 more locations