

Efficacy of EXPAREL vs. Bupivacaine

NCT07212114

Status	RECRUITING
Phase	Phase 3
Sponsor	Emory University
Enrollment	104 participants

Key Eligibility Criteria

Inclusion (4)

- Indicated to undergo unilateral total ankle arthroplasty
- American Society of Anesthesiologists (ASA) physical status 1, 2, or 3
- Able to provide informed consent, adhere to the study schedule, and complete all study assessments
- Body Mass Index (BMI) ≤ 18 and ≥ 40 kg/m²

Exclusion (14)

- Allergy, hypersensitivity, intolerance, or contraindication to any of the study medications for which an alternative is not named in the protocol (e.g., amide-type local anesthetics, opioids, bupivacaine HCl, NSAIDs)
- Concurrent painful physical condition that may require analgesic treatment (such as long-term, consistent use of opioids) in the post-dosing period for pain, and which, in the Investigator's opinion, may confound the post-dosing assessments
- History of, suspected, or known addiction to or abuse of illicit drug(s), prescription medicine(s), or alcohol within the past 2 years
- Uncontrolled anxiety, schizophrenia, or other psychiatric disorder that, in the opinion of the Investigator, could interfere with study assessments or compliance
- Currently pregnant, nursing, or planning to become pregnant during the study
- ... and 9 more (see full listing online)

Locations (1 total)

Emory University Orthopaedic Hospital Musculoskeletal Institute (MSK), Atlanta, Georgia, United States