

Prophylactic Intravenous Calcium Gluconate to Decrease Blood Loss at Time of Cesarean Delivery in Pregnant Patients at High Risk for Uterine Atony

NCT07217899

Status	RECRUITING
Phase	Phase 2
Sponsor	University of Michigan
Enrollment	140 participants

Key Eligibility Criteria

Inclusion (14)

- English speaking
 - Viable Intrauterine Pregnancy, Gestational Age e24 weeks
 - Labor Converted to Cesarean Delivery at High Risk for Atonic Bleeding Defined as: Any exposure to oxytocin infusion for labor augmentation or induction prior to cesarean delivery
 - Or Scheduled Cesarean Delivery at High Risk for Atonic Bleeding Defined as scheduled Cesarean-Section (CS) plus any one of:
 - \> 4 Prior deliveries
- ... and 9 more (see full listing online)

Exclusion (21)

- Non-English speaking
 - Antenatal suspicion for placenta accreta spectrum
 - History of allergic reaction to Calcium Gluconate
 - Patients with hypertensive disorder of pregnancy receiving Magnesium Sulfate for seizure prophylaxis
 - Underlying Renal Disease defined as Cr\>1.0
- ... and 16 more (see full listing online)

Locations (1 total)

University of Michigan, Ann Arbor, Michigan, United States