

Comparing Numbing Techniques in Mohs Micrographic Surgery

NCT07222241

Status	RECRUITING
Phase	Not Applicable
Sponsor	Abramson Cancer Center at Penn Medicine
Enrollment	150 participants

Key Eligibility Criteria

Inclusion (5)

- Age ≥18 years.
- Scheduled to undergo MMS.
- Lesions on the face or scalp, defined as the region superior to the mandibular margin anteriorly, superior to the external occipital protuberance posteriorly, and superior to the mastoid processes laterally.
- Tumor diameter plus anticipated stage 1 MMS margin ≥2 cm in diameter OR any tumor on the lip, nose, or eyelid.
- Ability to complete patient reported outcome measures in English.

Exclusion (8)

- Known allergy or contraindication to lidocaine or epinephrine.
- Chronic opioid use or pre existing pain disorders that may interfere with pain reporting.
- Presence of scar tissue at the anatomic site of local infiltration or nerve block that may alter anesthetic penetration.
- Concurrent multi site Mohs procedure.
- Signs of skin and soft tissue infection at the anatomic site of local infiltration or nerve block.

... and 3 more (see full listing online)

Locations (1 total)

Perelman Center for Advanced Medicine, Philadelphia, Pennsylvania, United States