

# Goat Milk-Derived Formula vs. Undiluted Goat Milk in Infants Unable to Exclusively Breastfeed: Growth and Biomarker Analysis

NCT07225153

---

Status	RECRUITING
Phase	Not Applicable
Sponsor	Aga Khan University
Enrollment	15 participants

## Key Eligibility Criteria

---

### Inclusion (6)

- Infants aged = 08 -10 weeks with WAZ better than -1.8. (NIPS & ICF, 2019) Age Sex WAZ Weight in Kilograms 2 Month Boys WAZ>-1.8 4-5 kg 2 Month Girls WAZ>-1.8 4-5 kg Table 3: WAZ scores
  - Living in Matiari District.
  - Who has access to goat milk in their households.
  - The Breastfeeding group: Women who exclusively breastfeeding and not giving any sort of formula or animal derived milk.
  - Intervention Group: Women who do not breastfeed at all or give goat milk to their infants (2-3 feeds per day).
- ... and 1 more (see full listing online)

### Exclusion (7)

- Infant birth weight < WAZ -1.8 i.e. weight-for-age based on established growth standards, such as the World Health Organization (WHO).
  - Birth deformities or disorders, such as genetic disorders, aerodigestive problems, or congenital anomalies.
  - Plan to migrate during the next six months.
  - We will exclude the women who exclusive breast feed and continue breast feeding for first 4 months.
  - If the child is enrolled or included in any other interventional trial.
- ... and 2 more (see full listing online)

## Locations (1 total)

---

MATIARI Research & Training Centre, Matiari, Sindh, Pakistan