

The Efficacy and Safety of Short Term S-ketamine Infusion as an Adjunctive Therapy in Patients With Fibromyalgia

NCT07230171

Status	RECRUITING
Phase	Not Applicable
Sponsor	Beijing Tiantan Hospital
Enrollment	106 participants

Key Eligibility Criteria

Inclusion (4)

- Patients between ages over 18 years with proper cognitive function and language skills for the study;
- Patients diagnosed with FM as defined by the ACR FM diagnostic criteria;
- Patients who have experienced insufficient symptom relief with non-pharmacological treatments and have not previously received recommended pharmacological treatment for FM;
- A score of e 4 on the average pain intensity over 7 days on the numeric rating scale (NRS) at baseline.

Exclusion (12)

- Patient refusal;
- Inability to sign informed consent;
- Had other secondary FM, this is, hypothyroidism, nutritional deficiency, diabetes mellitus, connective tissue disorder;
- Had psychiatric disorder, this is, schizophrenia and other psychotic disorder, bipolar disorder, or personality disorder;
- Obstructive sleep apnea syndrome or a STOP-Bang score e 3;
- ... and 7 more (see full listing online)

Locations (1 total)

Beijing Tiantan Hospital, Beijing, Beijing 100070, Beijing, China