

Risks and Benefits of Radiofrequency Ablation for Chronic Venous Insufficiency (CEAP C3-C6) in Patients Aged 80 and Over

NCT07233616

Status RECRUITING
Sponsor Chengdu University of Traditional Chinese Medicine
Enrollment 50 participants

Key Eligibility Criteria

Inclusion (9)

- Age: Aged 80 years or older.
 - Clinical Diagnosis: Symptomatic chronic venous insufficiency of the lower extremity, with the highest clinical class (CEAP classification) being one of the following:
 - C3: Venous edema
 - C4a: Pigmentation or eczema
 - C4b: Lipodermatosclerosis or atrophie blanche
- ... and 4 more (see full listing online)

Exclusion (9)

- Participants who meet any of the following criteria will be excluded from the study:
 - Mobility: Non-ambulatory or bedridden.
 - Thrombosis: Presence of acute superficial thrombophlebitis of the target vein, or deep vein thrombosis (DVT) in the target limb.
 - Arterial Disease: Significant peripheral arterial disease, defined as an ankle-brachial index (ABI) < 0.8 in the target limb.
 - Coagulopathy: Uncorrectable coagulation disorder.
- ... and 4 more (see full listing online)

Locations (1 total)

Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, China