

Genotype and Disease Burden of RSV in Older Vietnamese Adults (RSV: Respiratory Syncytial Virus)

NCT07239583

Status	RECRUITING
Sponsor	Tam Anh Research Institute
Enrollment	1,000 participants

Key Eligibility Criteria

Inclusion (6)

- Individuals \geq 60 years old at the time of screening (calculated from the date of birth to the date of screening)
- ARI or exacerbation of cardiopulmonary diseases diagnosed at the time of admission. ARI or ARI-compatible symptoms were defined as either:
 - New onset or increase from baseline of: nasal congestion, rhinorrhea, sore throat, hoarseness, cough, sputum production, dyspnea, wheezing, hypoxemia, or
 - Admitting diagnosis suggestive of ARI (e.g., pneumonia, upper respiratory infection, bronchitis, influenza, cough, viral respiratory illness, respiratory distress, or respiratory failure), or
 - Admitting diagnosis was the exacerbation of an underlying cardiac or pulmonary disease involving acute respiratory symptoms within the previous 2 weeks (e.g., congestive heart failure [CHF], chronic obstructive pulmonary disease [COPD], or asthma exacerbation).

... and 1 more (see full listing online)

Exclusion (4)

- Symptom onset $>$ 6 days prior to screening date
- Acute symptoms confirmed to be unrelated to infection or as assessed by the investigator
- Admission to any hospital within the 30 days preceding the screening date
- Previously enrolled in the study in the last 45 days

Locations (1 total)

Tam Anh TP. Ho Chi Minh General Hospital, Ho Chi Minh City, Southern Vietnam, Vietnam