

MAPT Protocol: Fixation Versus Arthroplasty Surgical Treatments for Early Recovery After HIP Fracture (FASTER-HIP)

NCT07244211

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Southern California
Enrollment	600 participants

Key Eligibility Criteria

Inclusion (6)

- years of age or older undergoing surgery due to a minimally displaced femoral neck fracture
- The patient has a health condition affecting physical mobility.
- Complete fracture of the femoral neck (AO/OTA 31B) confirmed with anteroposterior and lateral hip radiographs, computed tomography, or magnetic resonance imaging.
- Minimally displaced fracture that could be, in the judgment of the attending surgeon, managed with either arthroplasty or in situ internal fixation without reduction.
- Low energy injury mechanism.

... and 1 more (see full listing online)

Exclusion (19)

- The patient is not clinically suitable for either compared treatment.
- Expected injury survival of less than 12 months.
- Terminal illness with expected survival of less than 12 months.
- Incarceration.
- Unable to obtain informed consent due to language barriers.

... and 14 more (see full listing online)

Locations (4 total)

University of California, Irvine, Irvine, California, United States
Los Angeles General Medical Center, Los Angeles, California, United States
University of Southern California, Los Angeles, California, United States
... and 1 more locations