

Tenecteplase Before Interhospital Transfer for EVT in Acute Anterior Circulation LVO at 4.5-24 Hours

NCT07253181

Status	RECRUITING
Phase	Phase 3
Sponsor	Xuanwu Hospital, Beijing
Enrollment	572 participants

Key Eligibility Criteria

Inclusion (10)

- Age of 18 years or older;
- AIS symptom onset to treatment initiation within 4.5 to 24 hours, stroke onset is defined as the time the patient was last known to be well (including wake-up stroke and unwitnessed stroke);
- Signs and symptoms consistent with the diagnosis of an acute anterior circulation ischemic stroke involving occlusion of the internal carotid artery (ICA), MCA (M1 or M2) vessels;
- Functionally independent (mRS 0-2) prior to stroke onset;
- Baseline National Institute of Health Stroke Scale (NIHSS) of 6-25;
- ... and 5 more (see full listing online)

Exclusion (18)

- Known hypersensitivity or allergy to any ingredients of Tenecteplase;
- Intended to receive IVT as standard-of-care therapy;
- Rapidly improving symptoms with NIHSS score ≤ 6 before randomization;
- Any contra-indication for IVT except for the time criterion;
- Known hereditary or acquired hemorrhagic diathesis;
- ... and 13 more (see full listing online)

Locations (2 total)

Xuanwu Hospital Capital Medical University, Beijing, Beijing Municipality, China
Affiliated Hospital of Shandong Second Medical University, Weifang, Shandong, China