

Early Versus Delayed Trial Without Catheter in Men With Acute Urinary Retention.

NCT07283484

Status	RECRUITING
Phase	Not Applicable
Sponsor	Liselot Ribbert
Enrollment	478 participants

Key Eligibility Criteria

Inclusion (4)

- Adult men (e18 years)
- Diagnosis of acute urinary retention (AUR) treated with a transurethral catheter (TUC) and alpha-blocker therapy (tamsulosin, silodosin, or alfuzosin)
- Mentally competent and able to understand the potential benefits and burdens of study participation
- Provision of written or digital informed consent

Exclusion (10)

- Failed prior TWOC within the preceding 30 days
- Initial urinary retention volume \geq 1500 mL
- Neurogenic bladder dysfunction (e.g., multiple sclerosis, spinal cord injury, spina bifida)
- History of prostate cancer with ISUP grade group e2
- History of active bladder cancer or ongoing surveillance for bladder cancer
- ... and 5 more (see full listing online)

Locations (11 total)

St. Jansdal, Harderwijk, Gelderland, Netherlands
Canisius Wilhelmina Hospital, Nijmegen, Gelderland, Netherlands
Zuyderland Ziekenhuis, Heerlen, Limburg, Netherlands
... and 8 more locations