

Ocular Proteomics Testing In Chronic Atrophy

NCT07298291

Status	RECRUITING
Phase	Not Applicable
Sponsor	ClinOmicsAI
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (5)

- Male or female subjects aged between 50 and 89 years, inclusive
- For the GA cohort: Clinical diagnosis of GA secondary to AMD or dAMD in the study eye, confirmed by fundus autofluorescence imaging and spectral-domain optical coherence tomography (SD-OCT).
- For the control cohort: No clinical evidence of retinal disease in either eye.
- Best-corrected visual acuity (BCVA) of e20/200 in the study eye, as measured by Early Treatment Diabetic Retinopathy Study (ETDRS) charts
- Ability and willingness to provide written informed consent before any study-specific procedures.

Exclusion (6)

- Presence of active ocular infection or inflammation in either eye at the time of screening.
- History of intraocular surgery in the study eye within the three months preceding enrollment.
- Intraocular pressure (IOP) greater than 21 mmHg in the study eye at screening
- Use of systemic anticoagulation therapy that cannot be safely discontinued prior to ACP.
- Pregnant or nursing women.

... and 1 more (see full listing online)

Locations (1 total)

Erie Retina Research, Erie, Pennsylvania, United States