

Posttraumatic Stress Disorder in Ectopic Pregnancy

NCT07306689

Status RECRUITING
Sponsor Jagiellonian University
Enrollment 200 participants

Plain Language Summary

This study examines the presence and impact of post-traumatic stress disorder (PTSD) in women who have been diagnosed with an ectopic pregnancy — a potentially dangerous situation where the fertilized egg implants outside the uterus, usually in the fallopian tube. The goal is to understand the psychological burden of this experience so that better emotional support and screening tools can be developed.

****You may be eligible if...****

- You have been diagnosed with an ectopic pregnancy
- Your pregnancy was 14 weeks or earlier at the time of diagnosis
- You are willing to participate in a psychological evaluation

****You may NOT be eligible if...****

- You have a heterotopic pregnancy — a rare condition where one pregnancy is inside the uterus and another is outside it at the same time

Talk to your doctor to see if this trial is right for you.

Key Eligibility Criteria

Inclusion (2)

- ectopic pregnancy
- gestational age up to 14 weeks.

Exclusion (1)

- heterotopic pregnancy

Locations (2 total)

Jagiellonian University Medical College, Department of Gynecology and Obstetrics, Krakow, Poland
Jagiellonian University Medical College, Krakow, Poland

<https://clinicaltrials.gov/study/NCT07306689>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).