

SAS 1: Develop Ocular Disease (OSD) Diagnostic, Algorithms, and Management Protocols for Cataract Surgery

NCT07323264

Status RECRUITING
Sponsor Louis Tong
Enrollment 110 participants

Key Eligibility Criteria

Inclusion (6)

- Symptomatic for dry eye based on DEQ5 (>6) and \wedge either TBUT less than 5 s or staining $>$ grade 1 in any zone of the cornea
- Either use of diquasfosal, cyclosporine eyedrops or tacrolimus eyedrops for dry eye disease in the last 2 weeks (and previously started for more than 2 months), OR current and visible lower punctal plugs in at least one eye
- DEQ5 ≤ 6 .
- Staining $<$ grade 1 in at least one corneal zone.
- Not using diquasfosal, cyclosporine, tacrolimus in last 2 months.

... and 1 more (see full listing online)

Exclusion (10)

- If there is another ocular condition which affects vision as much as cataract: active corneal stromal disease or scar or significant corneal endothelial disease
- Known diagnosis of active thyroid dysfunction, Sjogren disease and rheumatoid arthritis
- Intraocular surgery within the previous 6 months
- Laser-assisted in situ keratomileusis or pterygium surgery within the previous 3 months
- Central nervous system and hormonal drugs required within the last month and during the study

... and 5 more (see full listing online)

Locations (1 total)

Singapore Eye Research Institute, Singapore, Singapore