

# Single- vs Double-Layer Cesarean Scar Repair and Myometrial Thickness

NCT07323355

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Ankara City Hospital Bilkent
<b>Enrollment</b>	102 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- Female participants aged 18-44 years.
- Undergoing cesarean delivery at the study hospital's Obstetrics and Gynecology clinic.
- Evaluated as clinically eligible based on history, physical examination, abdominal ultrasonography, and routine preoperative laboratory tests, including:
- Complete blood count (CBC), biochemical profile, prothrombin time (PT), activated partial thromboplastin time (aPTT), blood group determination. No abnormalities detected in the above assessments.

### Exclusion (6)

- Presence of chronic diseases, including: Rheumatologic diseases, renal failure, vascular malformations, hypertension, cardiac disease, diabetes mellitus, obesity, thyroid disease, congenital hematologic disorders.
- Suspected or confirmed placenta previa or placenta accreta spectrum.
- Clinical chorioamnionitis.
- Anterior wall uterine myomas.
- History of low transverse nonstandard uterine incision.
- ... and 1 more (see full listing online)

## Locations (1 total)

---

Ankara Bilkent City Hospital, Ankara, Not Valid, Turkey (Türkiye)