

Effect of Titrated Administration of Ciprofol on Perioperative Hypotension in Elderly Patients Undergoing Laparoscopic Abdominal Surgery: A Randomized Controlled Trial

NCT07328958

Status	RECRUITING
Phase	Not Applicable
Sponsor	Sixth Affiliated Hospital, Sun Yat-sen University
Enrollment	500 participants

Key Eligibility Criteria

Inclusion (3)

- Age ≥65 years, scheduled for elective laparoscopic abdominal surgery
- American Society of Anesthesiologists (ASA) physical status I-III
- Requirement for invasive arterial blood pressure monitoring

Exclusion (9)

- Participation in other clinical trials that may interfere with the intervention or outcomes of this study
 - Severe hepatic or renal disease (GFR <30 mL/min/1.73 m², requirement for renal replacement therapy, or Child-Pugh class C liver function)
 - Uncontrolled severe hypertension (preoperative SBP ≥180 mmHg or DBP ≥110 mmHg)
 - Patients with severe mental disorders (such as schizophrenia), epilepsy, or Parkinson's disease, severe cognitive or intellectual impairment, severe visual or hearing impairments affecting assessment, or long-term alcohol abuse or use of sedative/analgesic medications;
 - Known allergy to drugs used in this study
- ... and 4 more (see full listing online)

Locations (2 total)

Sixth Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China
Sixth Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China