

Hard Tissue Augmentation With or Without Connective Tissue Graft in Immediate Esthetic Implants

NCT07333534

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Baghdad
Enrollment	28 participants

Key Eligibility Criteria

Inclusion (4)

- Patients with a non-restorable tooth or a remaining root without signs of acute infection in the maxillary esthetic zone.
- The failing tooth is an incisor, canine or first bicuspid in the maxilla bounded by natural sound teeth.
- Sufficient bone (>4 mm) apically and palatally to allow for proper implant positioning with sufficient primary stability (e35 N cm).
- Sufficient mesial-distal and interocclusal space for placement of the implant and definitive restoration.

Exclusion (6)

- Patients with systemic conditions affecting bone healing (e.g., uncontrolled diabetes).
- Teeth with current acute periapical infection.
- Dehiscence of the labial (facial) bone plate after extraction.
- Signs of uncontrolled periodontal disease.
- Heavy smokers and vulnerable groups (pregnant females and decision-impaired individuals).
- ... and 1 more (see full listing online)

Locations (1 total)

University of Baghdad/ College of Dentistry/ Department of Oral and Maxillofacial surgery/ Implant Unit, Baghdad, Rusafa, Iraq