

The Efficacy of Remimazolam on Incidence of Hypoxia During Sedated Transvaginal Oocyte Retrieval: A Multicenter Randomized Controlled Trial.

NCT07335536

Status	RECRUITING
Phase	Not Applicable
Sponsor	Diansan Su
Enrollment	400 participants

Key Eligibility Criteria

Inclusion (2)

- Subjects aged 18 to 50 years planning to undergo painless oocyte retrieval
- Subjects classified as American Society of Anesthesiologists (ASA) physical status I-II

Exclusion (6)

- Subjects with contraindications to general anesthesia, acute or severe cardiovascular disease history, acute upper respiratory infection, chronic obstructive pulmonary disease (COPD), asthma attacks, or uncontrolled hypertension
- Subjects with alcohol abuse
- Subjects with severe hepatic or renal insufficiency, severe cardiovascular disease, or psychiatric disorder history
- Subjects with hearing impairment preventing communication
- Subjects with a history of long-term sedative or opioid use, or allergies to benzodiazepines, flumazenil, opioids and their antidotes, propofol, eggs, or soy products

... and 1 more (see full listing online)

Locations (3 total)

Dongyang People's Hospital, Dongyang, Zhejiang, China

The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, China

Lishui People's Hospital, Lishui, Zhejiang, China