

Selective Total Mesorectal Excision Based on Intra-Operative Frozen Section From Local Excision in Rectal Cancer Undergoing Neoadjuvant Chemoradiotherapy

NCT07337811

Status	RECRUITING
Phase	Not Applicable
Sponsor	Sun Yat-sen University
Enrollment	27 participants

Key Eligibility Criteria

Inclusion (8)

- Pathologically confirmed rectal adenocarcinoma.
- Neoadjuvant therapy based on radiotherapy; post-radiotherapy evaluation e8 weeks later shows near clinical complete response (near-cCR) or partial response (PR) with residual tumor ≤ 2 cm.
- Tumor distance from the anal verge ≥ 5 cm.
- Age 18-75 years.
- No synchronous multiple primary cancers.
- ... and 3 more (see full listing online)

Exclusion (14)

- Age < 18 or > 75 years.
- Concurrent or previous malignancy within the past 5 years (except adequately treated basal-cell or squamous-cell skin carcinoma or carcinoma in situ of the cervix).
- Emergency surgery required for bowel obstruction, perforation, or bleeding.
- Prior colorectal surgery that may compromise intestinal reconstruction.
- Need for en-bloc multivisceral resection.
- ... and 9 more (see full listing online)

Locations (1 total)

Dept. of Colorectal Surgery, Sun Yat-sen University Cancer Center. Yuexiu District, Dongfeng East Road 651, Guangzhou, Guangong, China