

Flapless Periodontal Regeneration Using Hyaluronic Acid Versus Enamel Matrix Derivative

NCT07339605

Status	RECRUITING
Phase	Not Applicable
Sponsor	University of Valencia
Enrollment	60 participants

Key Eligibility Criteria

Inclusion (12)

- Periodontitis stage III or IV, grade A or B (Tonetti et al., 2017): attachment loss \geq 5mm, radiographic bone loss extending to the middle third and beyond, tooth loss \geq 4 due to periodontitis, probing depth (PD) \geq 6mm.
 - Older adults \geq 18 years. The number per sex shall be balanced.
 - Plaque index (PI) \leq 1 following initial periodontal therapy and hygiene instructions.
 - Bleeding on probing (BoP) \leq 10%.
 - Only patients with optimal compliance, assessed during etiological therapy, will be selected.
- ... and 7 more (see full listing online)

Exclusion (9)

- The participant is pregnant, breastfeeding or plans to become pregnant in the next 6 months.
 - Smoking \geq 10 cigarettes/day.
 - Daily alcohol intake $>$ 4U.
 - Chronic illness or reduced mental capacity that may influence compliance with the protocol.
 - Medications or drugs that alter the patient's healing or with concomitant oral manifestations that, in the opinion of the investigator, could interfere with the assessment of safety or efficacy.
- ... and 4 more (see full listing online)

Locations (1 total)

Clínica Odontológica de la Universidad de Valencia, Valencia, Valencia, Spain

<https://clinicaltrials.gov/study/NCT07339605>

DISCLAIMER: This document is for informational purposes only and does not constitute medical advice. Always consult your healthcare provider before enrolling in any clinical trial. Information may not be up to date — verify details at [ClinicalTrials.gov](https://clinicaltrials.gov). Generated by [ClinicalTrialsFinder.org](https://clinicaltrialsfinder.org).