

Personalisation of Mean Arterial Pressure in Adult Patients With Cardiogenic Shock

NCT07345559

Status	RECRUITING
Phase	Not Applicable
Sponsor	CMC Ambroise Paré
Enrollment	406 participants

Key Eligibility Criteria

Inclusion (5)

- Aged e18 years
- Cardiogenic shock state, according to the consensus definition,
- SCAI (Society for Cardiovascular Angiography and Interventions) classification e C
- Consent from the patient or close relative / trusted person or emergency inclusion procedure
- Benefiting from ciary of a social security scheme

Exclusion (12)

- Catecholamine infusion for more than 24 consecutive hours;
- CVP \leq 5 mm Hg at inclusion;
- MAP \geq 70 mmHg at inclusion;
- Chronic kidney disease stage G4 (defined by an eGFR between 15-29 ml/min/1.73 m²) or G5 (defined by an eGFR less than 15 ml/min/1.73 m²) according to the KDIGO CKD classification at inclusion;
- Chronic dialysis or presence of renal replacement therapy criteria at inclusion ;
- ... and 7 more (see full listing online)

Locations (11 total)

CHU d'Amiens-Picardie, Amiens, France
Hôpital Henri Mondor, Créteil, France
Hôpital Privé Jacques Cartier, Massy, France
... and 8 more locations